

# Package ‘rsmatrix’

March 15, 2022

**Title** Matrices for Repeat-Sales Price Indexes

**Version** 0.2.3

**Description** A small package for calculating the matrices in Shiller (1991, <[doi:10.1016/S1051-1377\(05\)80028-2](https://doi.org/10.1016/S1051-1377(05)80028-2)>) that serve as the foundation for many repeat-sales price indexes.

**Depends** R (>= 3.6)

**Imports** Matrix (>= 1.2-17), methods, stats

**License** MIT + file LICENSE

**Encoding** UTF-8

**URL** <https://github.com/marberts/rsmatrix>

**NeedsCompilation** no

**Author** Steve Martin [aut, cre, cph] (<<https://orcid.org/0000-0003-2544-9480>>)

**Maintainer** Steve Martin <[stevemartin041@gmail.com](mailto:stevemartin041@gmail.com)>

**Repository** CRAN

**Date/Publication** 2022-03-15 14:30:02 UTC

## R topics documented:

rsmatrix-package	2
rs_matrix	3
rs_pairs	4
rs_var	5
<b>Index</b>	<b>8</b>

---

rsmatrix-package

*Matrices for Repeat-Sales Price Indexes*

---

## Description

A small package for calculating the matrices in Shiller (1991) that serve as the foundation for many repeat-sales price indexes.

## Usage

Most repeat-sales price indexes used in practice require the matrices in Shiller (1991, sections I-II), e.g., S&P's Case-Shiller index, Teranet-National Bank's HPI, and formerly Statistics Canada's RPPI. The `rs_matrix()` function produces a function to easily construct these matrices. In most cases data need to be structured as sales pairs, which can be done with the `rs_pairs()` function.

## Contribution

The **McSpatial** package (formerly on CRAN) has some functionality for making repeat-sales indices. The functions in this package build off of those in the **rsi** package in Kirby-McGregor and Martin (2019), which also gives a good background on the theory of repeat-sales indexes.

## Author(s)

**Maintainer:** Steve Martin <stevemartin041@gmail.com>

## References

ILO, IMF, OECD, UN, World Bank, Eurostat. (2013). *Handbook on Residential Property Prices Indices (RPPIs)*. Eurostat.

Kirby-McGregor, M., and Martin, S. (2019). An R package for calculating repeat-sale price indices. *Romanian Statistical Review*, 3:17-33.

Shiller, R. J. (1991). Arithmetic repeat sales price estimators. *Journal of Housing Economics*, 1(1):110-126.

## See Also

<https://github.com/marberts/rsmatrix>

---

`rs_matrix`*Shiller's repeat-sales matrices*

---

### Description

Create a function to compute the  $Z$ ,  $X$ ,  $y$ , and  $Y$  matrices in Shiller (1991, sections I-II) from sales-pair data in order to calculate a repeat-sales price index.

### Usage

```
rs_matrix(t2, t1, p2, p1, f = NULL, sparse = FALSE)
```

### Arguments

<code>t2, t1</code>	A pair of vectors giving the time period of the second and first sale, respectively. Usually a vector of dates, but other values are possible if they can be coerced to character and sorted in chronological order (i.e., with <code>order()</code> ).
<code>p2, p1</code>	A pair of numeric vectors giving the price of the second and first sale, respectively.
<code>f</code>	An optional factor the same length as <code>t1</code> and <code>t2</code> , or a vector to be turned into a factor, that is used to group sales.
<code>sparse</code>	Should sparse matrices from the <b>Matrix</b> package be used (faster for large datasets), or regular dense matrices (the default)?

### Details

The function returned by `rs_matrix()` computes a generalization of the matrices in Shiller (1991, sections I-II) that are applicable to grouped data. These are useful for calculating separate indexes for many, say, cities without needing an explicit loop.

The  $Z$ ,  $X$ , and  $Y$  matrices are not well defined if either `t1` or `t2` have missing values, and an error is thrown in this case. Similarly, it should always be the case that `t2 > t1`, otherwise a warning is given.

### Value

A function that takes a single argument naming the desired matrix. It returns one of two matrices ( $Z$  and  $X$ ) or two vectors ( $y$  and  $Y$ ), either regular matrices if `sparse = FALSE`, or sparse matrices of class `dgCMatrix` if `sparse = TRUE`.

### References

- Bailey, M. J., Muth, R. F., and Nourse, H. O. (1963). A regression method for real estate price index construction. *Journal of the American Statistical Association*, 53(304):933-942.
- Shiller, R. J. (1991). Arithmetic repeat sales price estimators. *Journal of Housing Economics*, 1(1):110-126.

**See Also**

[rs\\_pairs](#) for turning sales data into sales pairs.

**Examples**

```
# Make some data
x <- data.frame(date = c(3, 2, 3, 2, 3, 3),
                date_prev = c(1, 1, 2, 1, 2, 1),
                price = 6:1,
                price_prev = 1)

# Calculate matrices
mat <- with(x, rs_matrix(date, date_prev, price, price_prev))
Z <- mat("Z") # Z matrix
X <- mat("X") # X matrix
y <- mat("y") # y vector
Y <- mat("Y") # Y vector

# Calculate the GRS index in Bailey, Muth, and Nourse (1963)
b <- solve(crossprod(Z), crossprod(Z, y))[, 1]
# or b <- qr.coef(qr(Z), y)
(grs <- exp(b) * 100)

# Standard errors
vcov <- rs_var(y - Z %*% b, Z)
sqrt(diag(vcov)) * grs # delta method

# Calculate the ARS index in Shiller (1991)
b <- solve(crossprod(Z, X), crossprod(Z, Y))[, 1]
# or b <- qr.coef(qr(crossprod(Z, X)), crossprod(Z, Y))
(ars <- 100 / b)

# Standard errors
vcov <- rs_var(Y - X %*% b, Z, X)
sqrt(diag(vcov)) * ars^2 / 100 # delta method

# Works with grouped data
x <- data.frame(date = c(3, 2, 3, 2),
                date_prev = c(2, 1, 2, 1),
                price = 4:1,
                price_prev = 1,
                group = c("a", "a", "b", "b"))

mat <- with(x, rs_matrix(date, date_prev, price, price_prev, group))
b <- solve(crossprod(mat("Z"), mat("X")), crossprod(mat("Z"), mat("Y")))[, 1]
100 / b
```

**Description**

Turn repeat-sales data into sales pairs that are suitable for making repeat-sales matrices.

**Usage**

```
rs_pairs(period, product)
```

**Arguments**

**period** A vector that gives the time period for each sale. Usually a vector of dates, but other values are possible if they can be sorted in chronological order (i.e., with `order()`).

**product** A vector (usually a factor) that gives the product identifier for each sale.

**Value**

A numeric vector of indices giving the position of the previous sale for each product, with the convention that the previous sale for the first sale is itself. The first position is returned in the case of ties.

**See Also**

[rs\\_matrix](#) for using sales pairs to make a repeat-sales index.

**Examples**

```
# Make sales pairs
x <- data.frame(id = c(1, 1, 1, 3, 2, 2, 3, 3),
               date = c(1, 2, 3, 2, 1, 3, 4, 1),
               price = c(1, 3, 2, 3, 1, 1, 1, 2))

pairs <- rs_pairs(x$date, x$id)

x[c("date_prev", "price_prev")] <- x[c("date", "price")][pairs, ]

x
```

---

rs\_var

*Robust variance matrix for repeat-sales indexes*


---

**Description**

Convenience function to compute a cluster-robust variance matrix for a linear regression, with or without instruments, where clustering occurs along one dimension. Useful for calculating a variance matrix when a regression is calculated manually.

**Usage**

```
rs_var(u, Z, X = Z, ids = seq_len(nrow(X)), df = NULL)
```

**Arguments**

u	An $n \times 1$ vector of residuals from a linear regression.
Z	An $n \times k$ matrix of instruments.
X	An $n \times k$ matrix of covariates.
ids	A factor of length $n$ , or something that can be coerced into one, that groups observations in u. By default each observation belongs to its own group.
df	An optional degrees of freedom correction. Default is Stata's small sample degrees of freedom correction.

**Details**

This function calculates the standard robust variance matrix for a linear regression, as in Manski (1988, section 8.1.2) or White (2001, Theorem 6.3); that is,  $(Z'X)^{-1}V(X'Z)^{-1}$ . It is useful when a regression is calculated by hand. This generalizes the variance matrix proposed by Shiller (1991, section II) when a property sells more than twice.

This function gives the same result as `vcovHC(x, type = 'sss', cluster = 'group')` from the **plm** package.

**Value**

A  $k \times k$  covariance matrix.

**References**

- Manski, C. (1988). *Analog Estimation Methods in Econometrics*. Chapman and Hall.
- Shiller, R. J. (1991). Arithmetic repeat sales price estimators. *Journal of Housing Economics*, 1(1):110-126.
- White, H. (2001). *Asymptotic Theory for Econometricians* (revised edition). Emerald Publishing.

**Examples**

```
# Makes some groups in mtcars
mtcars$clust<- letters[1:4]

# Matrices for regression
x <- model.matrix(~ cyl + disp, mtcars)
y <- matrix(mtcars$mpg)

# Regression coefficients
b <- solve(crossprod(x), crossprod(x, y))

# Residuals
r <- y - x %*% b

# Robust variance matrix
vcov <- rs_var(r, x, ids = mtcars$clust)

## Not run:
```

```
# Same as plm
library(plm)
mdl <- plm(mpg ~ cyl + disp, mtcars, model = 'pooling', index = 'clust')
vcov2 <- vcovHC(mdl, type = 'sss', cluster = 'group')
vcov - vcov2

## End(Not run)
```

# Index

`order()`, [3](#), [5](#)

`rs_matrix`, [3](#), [5](#)

`rs_matrix()`, [2](#)

`rs_pairs`, [4](#), [4](#)

`rs_pairs()`, [2](#)

`rs_var`, [5](#)

`rsmatrix` (`rsmatrix`-package), [2](#)

`rsmatrix`-package, [2](#)